arXiv:1604.02693v1 [math.NT] 10 Apr 2016

A parametrised family of Mordell curves

Ajai Choudhry and Arman Shamsi Zargar

Abstract

An elliptic curve defined by an equation of the type $y^2 = x^3 + d$ is called a Mordell curve. We obtain a parametrised family of Mordell curves whose rank, in general, is at least three, and whose torsion group is $\mathbb{Z}/3\mathbb{Z}$.

Keywords: Mordell curves; rank of elliptic curves.

Mathematics Subject Classification 2010: 11D25, 11G05.

Ever since Fermat's assertion that the only solution in positive integers of the equation $y^2 = x^3 - 2$ is $(x, y) = (3, 5)$ [9], the diophantine equation,

$$y^2 = x^3 + d, \tag{1}$$

has been subjected to extensive investigations ([1], [2], [3], [4], [6], [7], [8], [10], [11, Chapter 26, pp. 238–254], [12], [16]). Eq. (1) has now been solved for all integer values of d with $|d| \leq 10^7$ [1].

The elliptic curve represented by Eq. (1) is known as a Mordell curve. Further, for various integer values of d, we know the structure of the torsion subgroup of the group of rational points on the Mordell curve [5, Theorem 5.3, p. 134]. It is noteworthy that whenever the integer d is a nonzero perfect square different from 1, the torsion subgroup is necessarily $\mathbb{Z}/3\mathbb{Z}$.

In this paper, we construct a parametrised family of Mordell curves defined by an equation of the type,

$$y^2 = x^3 + k^2. \tag{2}$$

We shall show that the rank of the elliptic curves belonging to this family is, in general, at least three. The torsion subgroup of all curves defined by the equation (2) is $\mathbb{Z}/3\mathbb{Z}$ with $(0,\, k)$ being a torsion point of order 3. Despite the considerable literature related to Eq. (1), it appears that such a parametrised family of curves has not been obtained earlier.

We will first solve the system of diophantine equations,

$$v_1^2 = u_1^3 + k^2, \tag{3}$$

$$v_2^2 = u_2^3 + k^2, \tag{4}$$

$$v_3^2 = u_3^3 + k^2. \tag{5}$$

On writing,

$$u_1 = am, \quad u_2 = bm, \quad u_3 = cm, \tag{6}$$

and eliminating m first between Eq. (3) and Eq. (4), and then between Eq. (3) and Eq. (5), we get the following two equations:

$$b^3(v_1^2 - k^2) = a^3(v_2^2 - k^2), \tag{7}$$

$$c^3(v_1^2 - k^2) = a^3(v_3^2 - k^2). \tag{8}$$

To solve equations (7) and (8), we write,

$$v_1 = w_1 t + k, \quad v_2 = w_2 t + k, \quad v_3 = w_3 t + k,$$

when each of the two equations (7) and (8) can be readily solved to get a nonzero solution for t. Equating these two values of t, we get the condition,

$$(b^3 w_1 - a^3 w_2)(c^3 w_1^2 - a^3 w_3^2) = (c^3 w_1 - a^3 w_3)(b^3 w_1^2 - a^3 w_2^2). \tag{9}$$

Now Eq. (9) is a homogeneous cubic equation in the variables w_1, w_2 and w_3 and it represents a cubic curve in the projective plane. Further, a rational point on this curve is easily seen to be $(w_1,\, w_2,\, w_3) = (a^3,\, b^3,\, c^3)$. The tangent to the cubic curve (9) at the aforementioned rational point

necessarily intersects the curve (9) at another rational point which is thus easily found, and is given by,

$$w_1 = a^3(b^3 + c^3 - a^3), \;\; w_2 = b^3(c^3 + a^3 - b^3), \;\; w_3 = c^3(a^3 + b^3 - c^3).$$

With these values of w_1, w_2, w_3, we obtain the following solution of the simultaneous equations (7) and (8):

$$\begin{aligned} v_1 &= -(3a^6 - 2a^3b^3 - 2a^3c^3 - b^6 + 2b^3c^3 - c^6)r, \\ v_2 &= (a^6 + 2a^3b^3 - 2a^3c^3 - 3b^6 + 2b^3c^3 + c^6)r, \\ v_3 &= (a^6 - 2a^3b^3 + 2a^3c^3 + b^6 + 2b^3c^3 - 3c^6)r, \\ k &= (a^6 - 2a^3b^3 - 2a^3c^3 + b^6 - 2b^3c^3 + c^6)r, \end{aligned} \tag{10}$$

where a, b, c and r are arbitrary parameters. Substituting the values of v_1, v_2, v_3 and k given by (10) and the value of u_1 given by (6) in (3), we get,

$$a^3m^3 = -8a^3r^2(a^3 + b^3 - c^3)(b^3 + c^3 - a^3)(c^3 + a^3 - b^3). \tag{11}$$

Now Eq. (11) is readily solved by taking $m = r$, when we get,

$$r = -8(a^3 + b^3 - c^3)(b^3 + c^3 - a^3)(c^3 + a^3 - b^3).$$

We thus obtain a solution of the system of equations (3), (4) and (5) which is given by,

$$\begin{aligned} k = -8&(a^3 + b^3 - c^3)(b^3 + c^3 - a^3)(c^3 + a^3 - b^3) \\ &\times (a^6 - 2a^3b^3 - 2a^3c^3 + b^6 - 2b^3c^3 + c^6), \end{aligned} \tag{12}$$

and

$$
\begin{aligned}
u_1 &= -8a(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3),\\
u_2 &= -8b(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3),\\
u_3 &= -8c(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3),\\
v_1 &= 8(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3)\\
&\qquad \times (3a^6-2a^3b^3-2a^3c^3-b^6+2b^3c^3-c^6),\\
v_2 &= -8(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3)\\
&\qquad \times (a^6+2a^3b^3-2a^3c^3-3b^6+2b^3c^3+c^6),\\
v_3 &= -8(a^3+b^3-c^3)(b^3+c^3-a^3)(c^3+a^3-b^3)\\
&\qquad \times (a^6-2a^3b^3+2a^3c^3+b^6+2b^3c^3-3c^6),
\end{aligned}
\tag{13}
$$

where a, b, c are arbitrary parameters.

It follows that when k is given by (12), there are three rational points $P_1(a,\, b,\, c)$, $P_2(a,\, b,\, c)$ and $P_3(a,\, b,\, c)$ on the elliptic curve (2) with co-ordinates $(u_i,\, v_i)$, $i = 1,\, 2,\, 3$, where the values of $u_i,\, v_i, i = 1,\, 2,\, 3$, are given by (13).

We will now apply a theorem of Silverman [15, Theorem 11.4, p. 271] to show that these points are linearly independent. For this, we must find a specialisation $(a, b, c) = (a_0, b_0, c_0)$ such that the points $P_1(a_0, b_0, c_0)$, $P_2(a_0, b_0, c_0)$, and $P_3(a_0, b_0, c_0)$ are linearly independent on the specialised curve over $\mathbb{Q}$.

We take $(a, b, c) = (1, 2, 3)$, when we get the elliptic curve,

$$
y^2 = x^3 + 28592640^2, \tag{14}
$$

on which we get the three points,

$$
\begin{aligned}
P_1(1,2,3) &= (97920, 41909760),\\
P_2(1,2,3) &= (195840, 91261440),\\
P_3(1,2,3) &= (293760, 161763840),
\end{aligned}
$$

each of which is of infinite order. The regulator of these three points, as determined by the software SAGE [13] is 33.9574760167017. As this is nonzero, it follows from a well-known theorem [14, Theorem 8.1, p. 242] that these three points are linearly independent. Hence the rank of the Mordell curve (14) is at least three.

It follows that, in general, the rank of the elliptic curves belonging to the parametrised family of Mordell curves (2), where k is given by (12), is at least three.

References

[1] M. Bennett and A. Ghadermarzi, Mordell's equation: a classical approach, *LMS J. Comput. Math.* **18**(1) (2015) 633–646.

[2] W. J. Ellison, F. Ellison, J. Pesek, C. E. Stahl and D. S. Stall, The diophantine equation $y^2 + k = x^3$, *J. Number Theory* **4**(2) (1972) 107–117.

[3] J. Gebel, A. Pethö and H. G. Zimmer, On Mordell's equation, *Compos. Math.* **110**(3) (1998) 335–367.

[4] M. Hall, Jr., Some equations $y^2 = x^3 - k$ without integer solutions, *J. Lond. Math. Soc. (1)* **28**(3) (1953) 379–383.

[5] A. Knapp, *Elliptic Curves* (Princeton University Press, 1992).

[6] M. Lal, M. F. Jones and W. J. Blundon, Numerical solutions of the diophantine equation $y^3 - x^2 = k$, *Math. Comp.* **20**(94) (1966) 322–325.

[7] W. Ljunggren, On the diophantine equation $y^2 - k = x^3$, *Acta Arith.* **8**(4) (1963) 451–463.

[8] L. J. Mordell, The diophantine equation $y^2 + k = x^3$, *Proc. Lond. Math. Soc. (2)* **13**(1) (1914) 60–80.

[9] L. J. Mordell, A statement by Fermat, *Proc. Lond. Math. Soc. (2)* **18**(1) (1920).

[10] L. J. Mordell, The infinity of rational solutions of $y^2 = x^3 + k$, *J. Lond. Math. Soc. (1)* **41**(1) (1966) 523–525.

[11] L. J. Mordell, *Diophantine equations* (Academic Press, London, 1969).

[12] D. Poulakis, The number of solutions of the Mordell equation, *Acta Arith.* **88**(2) (1999) 173–179.

[13] SAGE software, *Version 4.5.3*, http://www.sagemath.org.

[14] S. Schmitt and H. G. Zimmer, *Elliptic Curves: A Computational Approach* (Walter de Gruyter, Berlin, 2003).

[15] J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves* (Springer, New York, 1994).

[16] M. P. Young, The number of solutions to Mordell's equation in constrained ranges. *Mathematika* **61**(3) (2015) 708–718.

Postal Address 1: Ajai Choudhry,
13/4 A Clay Square,
Lucknow - 226001, INDIA.
E-mail: ajaic203@yahoo.com
Postal Address 2: Arman Shamsi Zargar,
Young Researchers and Elite Club, Ardabil Branch,
Islamic Azad University, Ardabil, Iran.
E-mail: shzargar.arman@gmail.com